arXiv:1304.3587v1 [math.DS] 12 Apr 2013

0-1 sequences of the Thue-Morse type and Sarnak's conjecture

El Houcein El Abdalaoui Stanisław Kasjan
Mariusz Lemańczyk*

April 15, 2013

Abstract

We show that the images via $z \mapsto z^m$ of the continuous part of the spectral measures of the dynamical systems generated by the 0-1 sequences of the Thue-Morse type are pairwise mutually singular for different odd numbers $m \in \mathbb{N}$. Sarnak's conjecture on orthogonality with the Möbius function is shown to hold for such dynamical systems.

1 Introduction

In this note, we consider dynamical systems arising from the $0-1$-sequences having the classical Thue-Morse sequence

$$x = (x(n)) = 0110100110010110\ldots$$

as a stabilizing subsequence (in what follows, we call such sequences of being of Thue-Morse type). More precisely, we consider (see Section 7 below) sequences $y \in \{0,1\}^{\mathbb{N}}$ having an increasing subsequence $1 \leq \ell_1 < \ell_2 < \ldots$ such that y arises from x by replacing its 0's and 1's at the first 2^{ℓ_n}-positions by a certain block $A_n \in \{0,1\}^{2^{\ell_n}}$ and $\widetilde{A}_n$ respectively[1], $n \geq 1$. Such $0-1$-sequences form a subclass of the family of *generalized Morse sequences* introduced by M. Keane [13] in 1968. We deal with the problem of the mutual singularity of the images via the maps $\mathbb{S}^1 \ni z \mapsto z^m \in \mathbb{S}^1$, $m \in \mathbb{N}$, of the continuous part of the spectral type of the corresponding dynamical system (Y, T, μ_y) [2]. Although the spectral theory of Morse dynamical systems is rather well-known [7], [14], the particular spectral problem formulated above seems not to be taken up yet in the literature. One of the mains results of the note shows that if $\widetilde{\sigma}$ denotes the continuous part of the spectral type of the dynamical system (Y, T, μ_y) arising from a Thue-Morse type sequence y, then $\widetilde{\sigma}^{(r)} \perp \widetilde{\sigma}^{(s)}$ whenever neither r is a 2^k-multiple (for some $k \geq 0$) of s, nor vice versa; here, for $m \in \mathbb{N}$, $\widetilde{\sigma}^{(m)}$ stands for the image of

---

*Research supported by Narodowe Centrum Nauki grant DEC-2011/03/B/ST1/00407

[1] $\widetilde{A}_n$ is the the block arising from A_n by interchanging 0's and 1's.

[2] $Y \subset \{0,1\}^{\mathbb{Z}}$ consists of all sequences $u \in \{0,1\}^{\mathbb{Z}}$ such that any block of consecutive symbols appearing in u also appears in y; T stands for the shift; the sequence y itself has an extension to a two-sided sequence in Y and, in fact, there are two such extensions which shows that the systems considered in this note are not distal, cf. [16].

$\widetilde{\sigma}$ via $z \mapsto z^m$. In particular, we have the mutual singularity of measures $\widetilde{\sigma}^{(r)}$ and $\widetilde{\sigma}^{(s)}$ whenever r, s are two different odd numbers.

Recently, such mutual singularity questions became of natural interest in connection with Sarnak's conjecture [18] which says that whenever (X, T) is a uniquely ergodic system[3] with zero (topological) entropy then

$$\frac{1}{N}\sum_{n=1}^{N} f(T^n u)\boldsymbol{\mu}(n) \to 0\ ^{4}, \tag{1}$$

where $\boldsymbol{\mu}(\cdot)$ stands for the classical Möbius function[5] and $u \in X$, $f \in C(X)$ are arbitrary. In numerous cases, this conjecture has recently been proved to hold: [1], [2], [3], [6], [16], [20]. For a connection of this conjecture with the classical Chowla conjecture see [18], [19] [6].

As noticed in [19], for the classical Thue-Morse sequence $x = (x(n)) = 0110100110010110\ldots$, it follows from [4] that

$$\frac{1}{N}\sum_{n=1}^{N} (-1)^{x(n)}\boldsymbol{\mu}(n) \to 0. \tag{2}$$

Another approach to handle (2) for x (as a matter of fact, to deal with so called q-multiplicative sequences - the Thue-Morse sequence is a 2-multiplicative sequence) proposed in [3] is to use the local dimension of the spectral measure of such a sequence and some recent results of Hochman [8]. Moreover, the orthogonality of the Thue-Morse sequence with the Möbius function also follows from [9] (see Theorem 1 therein) and the basic orthogonality criterion of Bourgain, Sarnak and Ziegler [3]. A natural question arises whether more than (2) is true, namely, whether Sarnak's conjecture holds for the associated Thue-Morse dynamical system, or more generally, for all Thue-Morse type dynamical systems. In the present note we provide the positive answer to the question and, moreover, it seems that we use the simplest available method. Indeed, the idea to use Furstenberg's disjointness to deal with Sarnak's conjecture for a uniquely ergodic system (X, T) has already appeared in [3]. Although, this idea cannot be always applied directly (for example, if the systems under consideration are not weakly mixing, which is the case of the present note), the spectral approach does help, see recent [1], [2]. This approach – a mixture of spectral disjointness with a direct proof of the Möbius orthogonality of sequences corresponding to the discrete part of the spectrum – will also be applied here.

The plan of the note is, firstly, to give a complete proof of the mutual spectral singularity result for the spectral measure of the classical Thue-Morse dynamical system (X, T, μ_x) itself via analysis of the Fourier transform of the spectral measure and using a simple general singularity criterion. This will allow us to reduce the problem of the validity of Sarnak's conjecture for (X, T, μ_x) to the same problem for some natural topological factor. This factor is determined by the so called Thue-Toeplitz sequence [7]. In order to complete the proof, we show that Sarnak's conjecture is true for all dynamical systems given by so called regular Toeplitz sequences [10], [21], see Section 6. This general result, in Section 7, enables us to conclude the validity of Sarnak's conjecture for all dynamical systems arising from the Thue-Morse type $0-1$-sequences.

---

[3] X here is a compact metric space and T is a homeomorphism of it.

[4] Following [18], when (1) holds, we speak about the orthogonality of the sequence $(f(T^n u))$ with the Möbius function.

[5] $\boldsymbol{\mu}(1) = 1$, $\boldsymbol{\mu}(n) = 0$ for any non-square free number n, and $\boldsymbol{\mu}(p_1 p_2 \ldots p_k) = (-1)^k$ for $p_1, p_2, \ldots, p_k$ different prime numbers.

[6] If the Chowla conjecture holds, also Sarnak's conjecture is valid.

[7] This factor is a particular uniquely ergodic model of the dyadic odometer.

Finally, we would like to warn the reader that even though our simple method of proving Sarnak's conjecture for the Thue-Morse dynamical system (and systems close to it) should work for many other systems of substitutional or Morse type, it cannot work for all generalized Morse sequences in view of a result of M. Guenais [7]. Indeed, in [7] it is shown that the problem whether there are Morse sequences for which the continuous part of the spectral measure of the corresponding dynamical system is Lebesgue is equivalent to the open classical problem of the existence of flat trigonometric polynomials with coefficients ± 1, i.e. we seek trigonometric polynomials $P_n(t) = \sum_{j=0}^{p_n-1} \epsilon_{j,n} e^{2\pi i j t}$ $(\epsilon_{j,n} = \pm 1)$ such that $\lim_{n\to\infty} \|P_n\|_{L^1}/\sqrt{p_n} = 1$.

For necessary notions and facts from ergodic theory and spectral theory of dynamical systems we refer the reader to e.g. [5], [12]. For the theory of substitutions, also from the dynamical system point of view, see [17].

2 Weighted operators and disjointness of spectral measures

Assume that S is an ergodic automorphism of a standard probability space $(Y, \mathcal{C}, \nu)$. Let $\varphi : Y \to \mathbb{Z}_2 := \{0, 1\}$ be a measurable map (cocycle). Denote by $W_{S,\varphi}$ the corresponding *weighted operator* on $L^2(Y, \mathcal{C}, \nu)$:

$$W_{S,\varphi}(\xi)(y) = (-1)^{\varphi(y)} \xi(Sy) \quad \text{for } \xi \in L^2(Y, \mathcal{C}, \nu). \tag{3}$$

Then, for each $m \in \mathbb{Z}$

$$\left(W^m_{S,\varphi}(\xi)\right)(y) = (-1)^{\varphi^{(m)}(y)} \xi(S^m y),$$

where $\varphi^{(m)}(y) = \varphi(y) + \varphi(Sy) + \ldots + \varphi(S^{m-1}y)$ for $m \in \mathbb{Z}$ positive and $\varphi^{(m_1+m_2)}(y) = \varphi^{(m_1)}(y) + \varphi^{(m_2)}(S^{m_1}y)$ for each $m_1, m_2 \in \mathbb{Z}$ $(\varphi^{(0)} = 0)$.

The following observation is folklore (we provide a proof for completeness).

Proposition 1. *Assume additionally that S has discrete spectrum. Assume moreover that $S^{q_n} \to Id$ on $L^2(Y, \mathcal{C}, \nu)$ for some increasing sequence $q_n \to \infty$ of natural numbers. If for some $c \in \mathbb{R}$, $|c| \le 1$*

$$\int_Y (-1)^{\varphi^{(q_n)}} \, d\nu \to c \tag{4}$$

then $(W_{S,\varphi})^{q_n} \to c \cdot Id$ weakly on $L^2(Y, \mathcal{C}, \nu)$.

Proof. By assumption, $L^2(Y, \mathcal{C}, \nu)$ is generated by eigenfunctions $\eta_i \in L^2(Y, \mathcal{C}, \nu)$ $(|\eta_i| = 1)$, $\eta_i \circ S = \alpha_i \cdot \eta_i$ for some $\alpha_i \in \mathbb{C}$ $(|\alpha_i| = 1)$, $i \ge 1$. Now, for $i, j \ge 1$ we have

$$\left\langle \, (W_{S,\varphi})^{q_n} \, \eta_i, \eta_j \right\rangle = \int_Y (-1)^{\varphi^{(q_n)}} \cdot \eta_i \circ S^{q_n} \cdot \overline{\eta}_j \, d\nu$$

$$= \alpha_i^{q_n} \int_Y (-1)^{\varphi^{(q_n)}} \cdot \eta_i \cdot \overline{\eta}_j \, d\nu \to c \langle \eta_i, \eta_j \rangle$$

since $\alpha_i^{q_n} \to 1$ and for $i \neq j$ we have

$$\begin{aligned}\int_Y (-1)^{\varphi^{(q_n)}} \cdot \eta_i \cdot \overline{\eta}_j \, d\nu &= \int_Y (-1)^{\varphi^{(q_n)}\circ S} \cdot (\eta_i \cdot \overline{\eta}_j) \circ S \, d\nu \\ &= \alpha_i \cdot \overline{\alpha}_j \int_Y (-1)^{\varphi^{(q_n)}\circ S - \varphi^{(q_n)}} (-1)^{\varphi^{(q_n)}} \cdot \eta_i \cdot \overline{\eta}_j \, d\nu \\ &= \alpha_i \cdot \overline{\alpha}_j \int_Y (-1)^{\varphi\circ S^{q_n} - \varphi} (-1)^{\varphi^{(q_n)}} \cdot \eta_i \cdot \overline{\eta}_j \, d\nu,\end{aligned}$$

so $\int_Y (-1)^{\varphi^{(q_n)}} \cdot \eta_i \cdot \overline{\eta}_j \, d\nu \to 0$ as (by the ergodicity of S) $\alpha_i \neq \alpha_j$ and $(-1)^{\varphi\circ S^{q_n} - \varphi} \to 1$ in measure. □

Given a measure σ on $\mathbb{S}^1$ denote by $\sigma^{(k)}$ the image of σ via the map $z \mapsto z^k$.

Consider now $\xi = \mathbf{1} = \mathbf{1}_Y$ and note that the Fourier transform of the spectral measure[8] $\sigma_{\mathbf{1}}$ of $\mathbf{1}$ is given by

$$\widehat{\sigma}_{\mathbf{1}}(m) = \int_Y (-1)^{\varphi^{(m)}} \, d\nu, \quad m \in \mathbb{Z}.$$

Since weak convergence in Proposition 1 is preserved when passing to closed, invariant subspaces we obtain the following result.

Corollary 1. *Under the assumptions of Proposition 1, if $r \neq s$ and for some $t \in \mathbb{Z}$ we have*

$$\int_Y (-1)^{\varphi^{(rtq_n)}} \, d\nu \to c_1, \tag{5}$$

$$\int_Y (-1)^{\varphi^{(stq_n)}} \, d\nu \to c_2 \tag{6}$$

with $c_1 \neq c_2$, then

$$\sigma_{\mathbf{1}}^{(r)} \perp \sigma_{\mathbf{1}}^{(s)}. \tag{7}$$

3 Generalized Morse sequences and strongly p-multiplicative sequences

Following [13], a sequence $x \in \{0,1\}^{\mathbb{N}}$ is called a *generalized Morse sequence* if

$$x = b^0 \times b^1 \times \ldots, \quad b^i \in \{0,1\}^{p_i}, b^i(0) = 0 \ (|b^i| := p_i \geq 2), \ i \geq 0,$$

where $B \times C = (B + C(0))(B + C(1)) \ldots (B + C(|C| - 1))$ with $B + 0 = B$ and $B + 1 =: \widetilde{B}$ arises from B by the interchange of 0s and 1s.

Consider now $X = \mathcal{O}(x) \subset \{0,1\}^{\mathbb{Z}}$ the subset of all two-sided sequences such that each block of consecutive symbols appearing in $y \in \mathcal{O}(x)$ also appears in x. This defines a closed, T-invariant (T is the shift by one position to the left) subset. Under some mild assumptions [13] on the blocks $b^0, b^1, \ldots$, one obtains a strictly ergodic dynamical system (X, μ_x, T), where μ_x is the unique T-invariant (Borel) probability measure (it is given by the average frequencies of blocks on x). From now on, only strictly ergodic case is considered.

---

[8] Given a unitary operator U of a Hilbert space $\mathcal{H}$ and $\xi \in \mathcal{H}$, the spectral measure σ_ξ is determined by $\widehat{\sigma}_\xi(m) := \int_{\mathbb{S}^1} z^m \, d\sigma_\xi(z) = \langle U^m \xi, \xi \rangle$ for all $m \in \mathbb{Z}$.

Consider the function $f : \mathcal{O}(x) \to \mathbb{R}$,

$$f(y) = (-1)^{y(0)}, \;\; y \in \mathcal{O}(x). \tag{8}$$

Then f is continuous, and the spectral measure σ_f of f under T is determined by [14], [15]

$$\widehat{\sigma}_f(m) = \int_{\mathcal{O}(x)} (-1)^{y(m)-y(0)}\, d\mu_x(y), \quad m \in \mathbb{Z}. \tag{9}$$

Moreover, the dynamical system $(\mathcal{O}(x), \mu_x, T)$ has a representation as a skew product S_φ over S which is the (q_n)-odometer ($q_n = p_0 \cdot \ldots \cdot p_{n-1}$, S is defined on $(Y, \mathcal{C}, \nu)$) and a cocycle $\varphi : Y \to \mathbb{Z}_2$ [13], [15] (called in [15] a Morse cocycle). From the spectral point of view, in this representation, the function f (given by (8)) corresponds to the function $\mathbf{1}_Y$ for the weighted operator $W_{S,\varphi}$, in other words [15]

$$\widehat{\sigma}_f(m) = \int_Y (-1)^{\varphi^{(m)}(y)}\, d\nu(y), \;\; m \in \mathbb{Z}. \tag{10}$$

A special case arises when $b^0 = b^1 = \ldots = B$. Then the corresponding (generalized) Morse sequence x can be obtained as a fixed point for the following substitution [17]

$$0 \mapsto B, \;\; 1 \mapsto \widetilde{B}.$$

In this particular case, the sequence $m_n(x) := (-1)^{x(n)}$, where $x = B \times B \times \ldots$, is *strongly p-multiplicative*, where p is the length of B, that is

$$m_{ap^n+b}(x) = m_a(x) \cdot m_b(x) \;\text{ for each } a, b \geq 0, \; b < p^n. \tag{11}$$

Each p-multiplicative sequence $(m_n(x))$ determines its *spectral measure* σ_x [17] whose Fourier transform $\widehat{\sigma}(\cdot)$ is given by

$$\widehat{\sigma}_x(k) = \lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{N} m_{n+k}(x) \cdot m_n(x), \;\; k \in \mathbb{Z}. \tag{12}$$

Since (by (11))

$$\widehat{\sigma}_x(kp) = \lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{N} m_{n+kp}(x) \cdot m_n(x) =$$

$$= \lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{[N/p]} \sum_{i=0}^{p-1} m_{pn+i+kp}(x) \cdot m_{pn+i}(x) =$$

$$\lim_{N\to\infty} \frac{1}{[N/p]} \sum_{n=1}^{[N/p]} m_{n+k}(x) \cdot m_n(x) \cdot m_i(x)^2 = \lim_{N\to\infty} \frac{1}{[N/p]} \sum_{n=1}^{[N/p]} m_{n+k}(x) \cdot m_n(x),$$

for each $k, n \geq 0$

$$\widehat{\sigma}_x(kp^n) = \widehat{\sigma}_x(k). \tag{13}$$

Corollary 2. *Assume that* $x = B \times B \times \ldots$ *Let* $r \neq s$. *If there exists* $t \geq 1$ *such that* $c_1 := \widehat{\sigma}(tr) \neq c_2 := \widehat{\sigma}(ts)$, *then* $\sigma_x^{(r)} \perp \sigma_x^{(s)}$.

Proof. It is well-known that $\sigma_f = \sigma_x$, e.g. [15]. The result then follows in view of Corollary 1 (with $q_n = p^n$), (10) and (13). $\square$

Remark 1. We can slightly strengthen this result by observing that whenever $|c_1| \neq |c_2|$ then $\sigma_x^{(r)} \perp \sigma_x^{(s)} * \delta_{z_0}$ for each $z_0 \in \mathbb{S}^1$. Indeed, we only need to show that whenever $z^{q_n} \to c$ weakly in $L^2(\mathbb{S}^1, \sigma)$ (that is, $\int_{\mathbb{S}^1} z^{q_n+m}\, d\sigma(z) \to c \int_{\mathbb{S}^1} z^m\, d\sigma(z)$ for each $m \in \mathbb{Z}$) and $z_0^{q_n} \to \alpha$, $|\alpha| = 1$, then $z^{q_n} \to \alpha c$ weakly in $L^2(\mathbb{S}^1, \sigma * \delta_{z_0})$.

Consider now $m_n = (-1)^{s_2(n)}$, where $s_2(n) = 0$ if the number of 1s in the binary expansion of n is even, and 1 otherwise. Directly from that we have $m_{2n} = m_n$ and $m_{2n+1} = -1$. Moreover, for each $n \geq 0$

$$m_n = m_n(01 \times 01 \times \ldots) \tag{14}$$

The sequence $x = 01 \times 01 \times \ldots$ is called the Thue-Morse sequence. From now on, we deal only with $\sigma = \sigma_x$, where x is the Morse-Thue sequence.

Since, for each $k \geq 0$, $\widehat{\sigma}(k) = \lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{N} m_{n+k} \cdot m_n$, we obtain

$$\widehat{\sigma}(2k) = \widehat{\sigma}(k), \;\; \widehat{\sigma}_{2k+1} = -\frac{1}{2}\big(\widehat{\sigma}(k) + \widehat{\sigma}(k+1)\big)\ {}^{9}. \tag{15}$$

Moreover, we can check directly that $\widehat{\sigma}(0) = 1$ and $\widehat{\sigma}(1) = -\frac{1}{3}$.

4 Fourier transform of the spectral measure of the Thue-Morse sequence. Spectral disjointness of powers

Following (15), we consider the sequence:

$$\begin{array}{ll} \widehat{\sigma}(0) = 1, \widehat{\sigma}(1) = -\frac{1}{3}, \widehat{\sigma}(2n) = \widehat{\sigma}(n), & \\ \widehat{\sigma}(2n+1) = -\frac{1}{2}(\widehat{\sigma}(n) + \widehat{\sigma}(n+1)) & \text{for} \quad n \geq 1. \end{array} \tag{16}$$

In what follows we need the values of $\widehat{\sigma}$ for small odd numbers[10]:

$$\begin{array}{llll} \widehat{\sigma}(1) = -\frac{1}{3}, & \widehat{\sigma}(3) = -\frac{1}{3}, & \widehat{\sigma}(5) = 0, & \widehat{\sigma}(7) = 0, \\ \widehat{\sigma}(9) = -\frac{1}{6}, & \widehat{\sigma}(11) = -\frac{1}{6}, & \widehat{\sigma}(13) = -\frac{1}{6}, & \widehat{\sigma}(15) = \frac{1}{6}, \end{array}$$

---

[9] Indeed, $m(1) = -1$, so

$$\frac{1}{N}\sum_{n=1}^{N} m_{n+2k+1} \cdot m_n =$$

$$\frac{1}{N}\left(\sum_{n=1}^{[N/2]} m_{2n+2k+1} \cdot m_{2n} + \sum_{n=1}^{[N/2]} m_{2n+1+2k+1} \cdot m_{2n+1} + \mathrm{O}(1)\right) =$$

$$-\frac{1}{2}\left(\frac{1}{[N/2]}\sum_{n=1}^{[N/2]} m_{n+k} \cdot m_n + \frac{1}{[N/2]}\sum_{n=1}^{[N/2]} m_{n+k+1} \cdot m_n\right) + \mathrm{o}(1/N),$$

whence (15) follows.

[10] We list some other values at prime instances to see that there are equalities between many Fourier coefficients: $\widehat{\sigma}(17) = \widehat{\sigma}(31) = \frac{1}{12}$, $\widehat{\sigma}(19) = \widehat{\sigma}(23) = \widehat{\sigma}(29) = -\frac{1}{12}$, $\widehat{\sigma}(37) = \widehat{\sigma}(41) = \widehat{\sigma}(59) = -\frac{1}{24}$, $\widehat{\sigma}(43) = \widehat{\sigma}(53) = \frac{1}{24}$, $\widehat{\sigma}(47) = \widehat{\sigma}(61) = -\frac{1}{8}$.

Lemma 1. *For any natural numbers* $n, a \geq 1$*, we have*

$$\widehat{\sigma}(2^a n + 1) = \left(-\frac{1}{2}\right)^a \left(\widehat{\sigma}(n+1) + \frac{1}{3}\widehat{\sigma}(n)\right) - \frac{1}{3}\widehat{\sigma}(n).$$

Proof. The assertion follows by induction on a. ☐

For a non-zero rational number $w = \frac{2^k p}{2^\ell q}$, where $p, q \in \mathbb{Z}$ are odd, by $v_2(w)$ we denote the multiplicity of 2 in w, i.e. $v_2(w) = k - l$.

Lemma 2. *If* $v_2(w_1) > v_2(w_2)$ *then* $v_2(w_1 + w_2) = v_2(w_2)$.

Proof. Writing $w_i = \frac{2^{k_i} p_i}{2^{l_i} q_i}$, $i = 1, 2$, we obtain

$$w_1 + w_2 = \frac{2^{k_2 + l_1}\left(2^{k_1 + l_2 - (k_2 + l_1)} p_1 q_2 + p_2 q_1\right)}{2^{l_1 + l_2} q_1 q_2}$$

since, by assumption, $k_1 + l_2 > k_2 + l_1$, and the result follows. ☐

Given an odd natural number K, we define sequences $K_0, \ldots, K_r$ and $a_1, \ldots, a_r$ of natural numbers, where each K_i is odd and:

$$K_0 = K,\ K_r = 1,$$
$$K_{i-1} = 2^{a_i} K_i + 1$$

for $i = 1, \ldots, r$. Set $r(K) := r$ and $l(K) := a_1 + \ldots + a_r$.

Lemma 3. *For an odd natural number* K*, we have* $l(K) = [\log_2 K]$*; in particular*

$$2^{l(K)} < K < 2^{l(K)+1}$$

Proof. Follows by induction on $r(K)$. ☐

Lemma 4. *If* K *is an odd natural number then* $\widehat{\sigma}(K) = 0$ *or* $v_2(\widehat{\sigma}(K)) \geq 2 - l(K)$*. If* $K \geq 9$ *then*

$$v_2(\widehat{\sigma}(K)) = 2 - l(K).$$

Proof. We proceed by induction on $l(K) \geq 3$. If $l(K) = 3$ then (see Lemma 3) $K \in \{9, 11, 13, 15\}$ and we see that the assertion is true.

Assume that $l(K) > 3$. Then $K = 2^{a_1} K_1 + 1$. We consider the cases $1 \leq K_1 \leq 7$ and $K_1 > 7$ separately.

If $K_1 = 1$ then $l(K) = a_1 > 3$ and (by Lemma 1)

$$\widehat{\sigma}(K) = (-\frac{1}{2})^{a_1} \left(\widehat{\sigma}(2) + \frac{1}{3}\widehat{\sigma}(1)\right) - \frac{1}{3}\widehat{\sigma}(1) = (-\frac{1}{2})^{a_1}(-\frac{4}{9}) + \frac{1}{9} = \frac{-(-1)^{a_1} + 2^{a_1 - 2}}{9 \cdot 2^{a_1 - 2}}$$

and $v_2(\widehat{\sigma}(K)) = 2 - a_1 = 2 - l(K)$.

If $K_1 = 3$ then $l(K) = a_1 + 1 > 3$ and

$$\widehat{\sigma}(K) = (-\frac{1}{2})^{a_1} \left(\widehat{\sigma}(4) + \frac{1}{3}\widehat{\sigma}(3)\right) - \frac{1}{3}\widehat{\sigma}(3) = (-\frac{1}{2})^{a_1}(-\frac{2}{9}) - \frac{1}{9} = \frac{-(-1)^{a_1} - 2^{a_1 - 1}}{9 \cdot 2^{a_1 - 1}}$$

and $v_2(\widehat{\sigma}(K)) = 1 - a_1 = 2 - l(K)$.

If $K_1 = 5$ then $l(K) = a_1 + 2 > 3$ and

$$\widehat{\sigma}(K) = \big(-\frac{1}{2}\big)^{a_1}\left(\widehat{\sigma}(6) + \frac{1}{3}\widehat{\sigma}(5)\right) - \frac{1}{3}\widehat{\sigma}(5) = \big(-\frac{1}{2}\big)^{a_1}\frac{1}{3} = \frac{(-1)^{a_1}}{3\cdot 2^{a_1}}$$

and $v_2(\widehat{\sigma}(K)) = -a_1 = 2 - l(K)$.

If $K_1 = 7$ then $l(K) = a_1 + 2 > 3$ and

$$\widehat{\sigma}(K) = \big(-\frac{1}{2}\big)^{a_1}\left(\widehat{\sigma}(8) + \frac{1}{3}\widehat{\sigma}(7)\right) - \frac{1}{3}\widehat{\sigma}(7) = \big(-\frac{1}{2}\big)^{a_1}\big(-\frac{1}{3}\big) = \frac{(-1)^{a_1+1}}{3\cdot 2^{a_1}}$$

and $v_2(\widehat{\sigma}(K)) = -a_1 = 2 - l(K)$.

Now consider the case $K_1 > 7$. Then $K_1 + 1 = 2(2^{a_2-1}K_2 + 1) = 2^s L$ for some $s \geq 1$ and an odd number L. Observe that $l(L) < l(K_1)$ (otherwise, $K_1 < 2^{m+1}$ and $2^m < L$ for some m which leads to a contradiction: $2^{m+s} < 2^s L = K_1 + 1 \leq 2^{m+1}$). It follows that either $\widehat{\sigma}(K_1 + 1) = \widehat{\sigma}(L) = 0$ or $v_2(\widehat{\sigma}(K_1 + 1)) = v_2(\widehat{\sigma}(L)) = 2 - l(L) > 2 - l(K_1)$ thanks to the inductive assumption. It follows that

$$v_2\left(\big(-\frac{1}{2}\big)^{a_1}\widehat{\sigma}(K_1 + 1)\right) > 2 - a_1 - l(K_1) = 2 - l(K),$$

and similarly

$$v_2\left(\big(-\frac{1}{2}\big)^{a_1}\frac{1}{3}\widehat{\sigma}(K_1)\right) = 2 - a_1 - l(K_1) = 2 - l(K)$$

by the inductive assumption. Moreover

$$v_2\left(\frac{1}{3}\widehat{\sigma}(K_1)\right) = 2 - l(K_1) > 2 - l(K).$$

Recalling that

$$\widehat{\sigma}(K) = \big(-\frac{1}{2}\big)^{a_1}\left(\widehat{\sigma}(K_1 + 1) + \frac{1}{3}\widehat{\sigma}(K_1)\right) - \frac{1}{3}\widehat{\sigma}(K_1),$$

using Lemma 2, we finally obtain that $v_2(\widehat{\sigma}(K)) = 2 - l(K)$. $\square$

Two numbers $K, L \geq 0$ are called *TM-equivalent* if $\widehat{\sigma}(2K + 1) = \widehat{\sigma}(2L + 1)$.

Lemma 5. *Assume that $K, L \geq 9$ are odd natural numbers. If L is TM-equivalent to K then $l(K) = l(L)$.*

Proof. Since $v_2(\widehat{\sigma}(K)) = v_2(\widehat{\sigma}(L)))$ and $K, L \geq 9$ then, in view of Lemma 4, we get $l(K) = l(L)$. $\square$

Finally, we need the following observation.

Lemma 6. *Assume that $r, s \geq 1$, $r < s$, are odd numbers. Then for any integer $a \geq 1$ large enough there exist an odd number t such that*

$$rt < 2^a < st.$$

Proof. It is enough to consider $s = r+2$. Then, for each $a \geq 1$ sufficiently large, we have

$$r\left(\frac{2^a}{r+1}+2\right) < 2^a < (r+2)\left(\frac{2^a}{r+1}-1\right),$$

so it is enough to take for t either $\left[\frac{2^a}{r+1}\right]$ or $\left[\frac{2^a}{r+1}\right]+1$ (for a sufficiently large). □

Using Lemmas 4, 6, 5 and Corollary 2 and noticing that $\sigma^{(k)} = \sigma^{(2k)}$ for each $k \geq 1$ (equivalently, σ is invariant under the map $z \mapsto z^2$), we obtain the following result.

Proposition 2. *Let σ denote the spectral measure associated to the Thue-Morse sequence $x = 01 \times 01 \times \ldots$ Then $\sigma^{(r)} \perp \sigma^{(s)}$ if and only if $\max\{r/s, s/r\} \notin \{2^a : a \in \mathbb{N}\}$. Moreover, $\sigma^{(r)} = \sigma^{(2^a r)}$ for each integers $a, r \geq 0$.*

Corollary 3. *Let σ denote the spectral measure associated to the Thue-Morse sequence $x = 01 \times 01 \times \ldots$ Then $\sigma^{(p)} \perp \sigma^{(q)}$ for arbitrary odd numbers $p \neq q$.*

Remark 2. Recall [17] that the maximal spectral type of the Thue-Morse dynamical system is given by the measure which is the sum of two measures: the discrete measure concentrated on all roots of unity of degree 2^n, $n \geq 0$, and the continuous measure $\widetilde{\sigma}$ which is the convolution of the discrete measure with σ. In view of Remark 1, we obtain that the assertions of Proposition 2 and Corollary 3 are true when σ is replaced by $\widetilde{\sigma}$.

5 On Sarnak's conjecture for the Thue-Morse dynamical system

As before, we let x denote the Thue-Morse sequence. Set $X = \mathcal{O}(x)$ and notice that the map τ which interchange 0s and 1s is a homeomorphism (involution) preserving X and commuting with the shift T. Denote also by x any extension of x to a two-sided member of X. Take $f \in C(X)$ arbitrary and note that

$$f = f_t + f_m, \tag{17}$$

where $f_t = \frac{1}{2}(f + f \circ \tau)$ and $f_m = \frac{1}{2}(f - f \circ \tau)$. Then $f_t = f_t \circ \tau$, $f_m \circ \tau = -f_m$.

Basic spectral theory for group extensions shows that the spectral measure σ_{f_m} of f_m is absolutely continuous with respect to $\widetilde{\sigma}$, so by Remark 2 and an observation from [1],

$$\frac{1}{N}\sum_{n=1}^{N} f_m(T^n y)\boldsymbol{\mu}(n) \to 0$$

for **each** $y \in \mathcal{O}(x)$ [11].

It follows that to show Sarnak's conjecture for the Thue-Morse dynamical system we need to prove it for the associated Toeplitz dynamical system [15]. We recall that the Toeplitz sequence z, called from now on, Thue-Topelitz, associated with the Thue-Morse sequence x arises as follows: At first step we we put $z_0 = z_2 = \ldots = z_{2n} = \ldots = 1$ and leave odd places undefined. At the second step we put $z_1 = z_5 = \ldots = z_{4n+1} = \ldots = 0$, that is, we fill every second unfilled place by putting 0 here. At the third step we repeat step one setting 1 at every second unfilled place, etc. It is not hard to see that for each $n \geq 0$

---

[11] In particular, it holds for $f(y) = (-1)^{y(0)}$ which shows that that not only x but each $y \in \mathcal{O}(x)$ is orthogonal to the Möbius function.

$$z = B_n?B_n?B_n?\ldots, \tag{18}$$

where $|B_n| = 2^n - 1$ [12], and "?" stand for un unfilled place (half of these unfilled places will be filled at step $n+1$ of the construction). To prove Sarnak's conjecture for $(\mathcal{O}(z), T)$, we need to consider a linearly dense set of functions in $C(\mathcal{O}(z))$.

Given a sequence $w = (w_i)_{i\in\mathbb{Z}}$ and $a \in \mathbb{Z}$, $l \geq 0$ we denote by $w[a, a+l)$ the sequence $(w_a, w_{a+1}, ..., w_{a+l-1})$. We consider maps defined as follows: Fix $\ell \geq 1$, $a \in \mathbb{Z}$ and consider $f : \{0,1\}^\ell \to \mathbb{C}$. It has a natural extension to a continuous map $f : \mathcal{O}(z) \to \mathbb{C}$ by setting

$$f(w) = f(w[a, a+\ell)), \quad w \in \mathcal{O}(z).$$

Under this notation, $f(T^k w) = f(w[a+k, a+k+\ell))$.

Observe that any continuous map on $\mathcal{O}(z)$ having only finitely many values is obtained this way.

Given $\ell \geq 1$, $a \in \mathbb{Z}$ and a sequence $w' \in \{0,1\}^\ell$, we denote by $U_{w',a}$ the open set $\{w \in \mathcal{O}(z) : w[a, a+\ell) = w'\}$.

Lemma 7. *The functions $f : \mathcal{O}(z) \to \mathbb{C}$ taking finitely many values form a dense subset in $C(\mathcal{O}(z))$.*

Proof. Fix $\varepsilon > 0$ and let $f \in C(\mathcal{O}(z))$. Since f is uniformly continuous, there exist $\ell \geq 1$ and $a < 0$ such that the diameter of $f(U_{w',a})$ is less than ε for any $w' \in \{0,1\}^\ell$. Fix such an $a \in \mathbb{Z}$. Let $w_1, \ldots, w_M \in \mathcal{O}(z)$ be the representatives of the equivalence classes of the relation $\sim$ defined by $w \sim w'$ if and only if $w[a, a+\ell) = w'[a, a+\ell)$. Let f' be given by the formula $f'(w) = f(w_j)$, where $w[a, a+\ell) = w_j[a, a+\ell)$. Then $|f(w) - f'(w)| < \varepsilon$ for any $w \in \mathcal{O}(z)$. □

Lemma 8. *We have $\frac{1}{N}\sum_{k=1}^N f(T^k w)\boldsymbol{\mu}(k) \to 0$ for any $f \in C(\mathcal{O}(z))$ taking finitely many values and any $w \in \mathcal{O}(z)$.*

Proof. We can assume that there are $a \in \mathbb{Z}$, $\ell \in \mathbb{N}$ such that the value $f(w)$ depends only on $w[a, a+\ell)$.

The map f is bounded, say $|f(w)| \leq F$ for all $w \in \mathcal{O}(z)$.

Fix $\varepsilon > 0$ and let n be a natural number such that

$$\frac{\ell F}{2^n} < \frac{\varepsilon}{3}. \tag{19}$$

We recall that $\frac{1}{N}\sum_{k=1}^N b_k\boldsymbol{\mu}(k) \to 0$ for any periodic (or eventually periodic with a fixed length of preperiod) sequence $(b_k) \subset \mathbb{C}$ (by the distribution of prime numbers in arithmetic progressions and the fact that the linear space of such sequences is of finite dimension).

Let M_1 be a number such that

$$\frac{1}{N}\sum_{k=1}^N b_k\boldsymbol{\mu}(k) < \frac{\varepsilon}{3 \cdot 2^n} \tag{20}$$

---

[12] In fact, $z(i) = x(i) + x(i+1) \bmod 2$.

for any periodic sequence (b_k) bounded by F of period at most 2^n and any $N \geq 2^n M_1$. Assume moreover that $\frac{F}{M_1} < \frac{\varepsilon}{3}$.

Take $N > 2^n M_1$, put $M = [N/2^n]$ and let $w \in \mathcal{O}(z)$. Then $w[1, N+1) = z[i, i+N)$ for some $i \in \mathbb{Z}$, thus (see (18))

$$w[1, N+1) = C' B_n y_1 B_n y_2 \dots B_n y_M C'',$$

where B_n is a $\{0,1\}$-sequence of length $2^n - 1$, C', C'' are $\{0,1\}$-sequences such that C' is a suffix of $B_n y_0$, C'' is a prefix of B_n and $y_1, \dots, y_M \in \{0,1\}$. Let c', c'' denote the length of C', C'', respectively.

Then $\sum_{k=1}^{N} f(T^k w)\boldsymbol{\mu}(k)$ equals

$$\begin{aligned}
&f(Tw)\boldsymbol{\mu}(1) + f(T^2 w)\boldsymbol{\mu}(2) + \ldots + f(T^{c'} w)\boldsymbol{\mu}(c') + \\
&f(T^{c'+1} w)\boldsymbol{\mu}(c'+1) + f(T^{c'+2^n+1} w)\boldsymbol{\mu}(c'+2^n+1) + \ldots + f(T^{c'+(M-1)2^n+1} w)\boldsymbol{\mu}(c'+(M-1)2^n+1) + \\
&f(T^{c'+2} w)\boldsymbol{\mu}(c'+2) + f(T^{c'+2^n+2} w)\boldsymbol{\mu}(c'+2^n+2) + \ldots + f(T^{c'+(M-1)2^n+2} w)\boldsymbol{\mu}(c'+(M-1)2^n+2) + \\
&\ldots\ldots \\
&f(T^{c'+2^n} w)\boldsymbol{\mu}(c'+2^n) + f(T^{c'+2\cdot 2^n} w)\boldsymbol{\mu}(c'+2\cdot 2^n) + \ldots + f(T^{c'+M2^n} w)\boldsymbol{\mu}(c'+M2^n) + \\
&f(T^{c'+M2^n+1} w)\boldsymbol{\mu}(c'+M2^n+1) + f(T^{c'+M2^n+2} w)\boldsymbol{\mu}(c'+M2^n+2) + \ldots + f(T^{c'+M2^n+c''} w)\boldsymbol{\mu}(c'+M2^n+c'')
\end{aligned}$$

Having organized the sum into $2^n + 2$ rows as above, we denote by E' the sum in the first row, by E'' the sum in the last one and by Σ_i the sum in the $i+1$-st row for $i = 1, \dots, 2^n$.

Observe that

$$|E' + E''| \leq 2^n F. \tag{21}$$

Moreover, Σ_i is an expression of the form $\Sigma_{k=1}^{N} b_k \boldsymbol{\mu}(k)$, where (b_k) is a periodic sequence,[13] provided the segment $w[c'+i, c'+i+\ell)$ does not meet any "uncertain" entry y_j. The latest condition holds for $i \leq 2^n - \ell$.

It follows by (20) that

$$|\Sigma_i| \leq \frac{N\varepsilon}{3 \cdot 2^n}$$

for $i = 1, \dots, 2^n - \ell$.

For the remaining i's we have

$$|\Sigma_i| \leq MF. \tag{22}$$

Finally, taking into account (19), (21), (22) and $\frac{F}{M_1} < \frac{\varepsilon}{3}$, we get

$$\left| \frac{1}{N} \sum_{k=1}^{N} f(T^k w)\boldsymbol{\mu}(k) \right| < \frac{2^n F}{N} + \frac{2^n \varepsilon}{3 \cdot 2^n} + \frac{MF}{N} \leq \varepsilon.$$

□

Proposition 3. $\frac{1}{N}\sum_{k=1}^{N} f(T^k w)\boldsymbol{\mu}(k) \to 0$ *for any* $f \in C(\mathcal{O}(z))$ *and any* $w \in \mathcal{O}(z)$.

Proof. Follows by Lemma 8 and Lemma 7 and the fact that $\frac{1}{N}\sum_{k=1}^{N} \boldsymbol{\mu}(k) \to 0$. □

---

[13] The periodic sequence (b_k) has form $(0, ..., 0, \phi, 0, ..., 0, \phi, 0, ...)$ and period 2^n, where ϕ is a fixed value of f.

6 On Sarnak's conjecture for regular Toeplitz dynamical systems

We will consider sequences of a finite alphabet $A=\{0,\ldots,d-1\}$, for some $d\geq 2$. Following [10], a sequence $z\in A^{\mathbb{N}}$ is called a Toeplitz sequence if for each $n\geq 0$ there is $a_n\geq 1$ such that $z(n)=z(n+a_n)=z(n+2a_n)=\ldots$ It can be proved [10], [21] that then there is an increasing sequence (p_n), $p_n|p_{n+1}$ such that for each $n\geq 1$, z can be represented as $z=C_nC_n\ldots$ with C_n being a block over the alphabet $A\cup\{?\}$, where the sign "?" means an "unfilled" place (at the nth step of the construction of z) and $|C_n|=p_n$, $n\geq 1$. Recall that a Toeplitz sequence z is called *regular* if

$$\text{(the number of unfilled places in } C_n)/p_n\to 0\ \text{ when } n\to\infty.$$

The dynamical system generated by a regular Toeplitz sequence is uniquely ergodic [10], [21]. Moreover, dynamical systems generated by Toeplitz sequences are so called almost 1-1 extensions of their maximal equicontinuous factors which are (p_n)-odometers.

Now, the same method which has been used to show that Sarnak's conjecture holds for the Thue-Toeplitz sequence shows the following.

Proposition 4. *Let z be any regular Toeplitz sequence. Then the dynamical systems $(\mathcal{O}(z),T)$ determined by z satisfies (1).*

7 Generalized Morse sequences having a stabilizing Thue-Morse subsequence

We now come back to the cocycle representation of Morse dynamical systems, to see how to obtain more systems for which Sarnak's conjecture holds.

Let $y=b^0\times b^1\times\ldots$ be a (generalized) Morse sequence. Fix $s\geq 1$ and then $k,K\geq 1$ (we think of K as going to ∞) and write

$$y=B\times C\times b^{k+K}\times b^{k+K+1}\times\ldots,\tag{23}$$

where $B=b^0\times\ldots\times b^{k-1}$, $C=b^k\times\ldots\times b^{k+K-1}$. Notice that this operation of putting parentheses does not change the space $\mathcal{O}(y)$, in particular, the number (cf. (9))

$$\int_{\mathcal{O}(y)}(-1)^{u(s|B|)-u(0)}\,d\mu_y(u)=\int_{\mathcal{O}(y)}(-1)^{u(sq_k)-u(0)}\,d\mu_y(u)=\widehat{\sigma}_y(sq_k)\tag{24}$$

is not changed compared to the original representation of y. However, introducing parentheses does change the cocycle representation of the corresponding dynamical system (the underlying odometer has a different algebraic representation, and the Morse cocycle also changes). We now provide some details (see [15]) concerning the first two steps in the definition of the odometer and the Morse cocycle.

To the length $|B|$ of $B:=(b_0,b_1,\ldots,b_{|B|-1})$ there corresponds a partition $\mathcal{D}^{(1)}=\{D_0^{(1)},\ldots,D_{|B|-1}^{(1)}\}$ for which $SD_i^{(1)}=D_{i+1}^{(1)}$ modulo $|B|$. At this stage, the Morse cocycle $\phi : Y \to \{0,1\}$ is defined as being constant on each $D_i^{(1)}$, $i = 0, 1, \ldots, |B| - 2$ with

$$\phi|_{D_i^{(1)}} = b_i + b_{i+1} \text{ (modulo 2)}, \ i = 0, 1, \ldots, |B| - 2,$$

i.e. we set the successive values of the block $\check{B} := (b_0 + b_1)(b_1 + b_2) \ldots (b_{|B|-2} + b_{|B|-1})$ as the values of ϕ on $D_0^{(1)}, \ldots, D_{|B|-2}^{(1)}$, and ϕ is not defined on $D_{|B|-1}^{(1)}$ (modulo 2). Also, denote $\Sigma := \check{b}_0 + \ldots + \check{b}_{|B|-2}$. At the next step, the tower $\mathcal{D}^{(1)}$ is refined to a tower $\mathcal{D}^{(2)}$ in the following way: We divide the base $D_0^{(1)}$ into $|C|$ equal pieces, which yields the partition of $\mathcal{D}^{(1)}$ into columns and obtain a new tower $\mathcal{D}^{(2)}$ (with $|B| \cdot |C|$ levels) in which

$$SD_i^{(2)} = D_{i+1}^{(2)} \text{ modulo } |B| \cdot |C| \text{ and } D_{|B|-1}^{(1)} = \bigcup_{j=1}^{|C|} D_{j|B|-1}^{(2)}.$$

We need to define ϕ on the levels of $\mathcal{D}^{(2)}$ (a constant values on each level) on which ϕ is not defined yet, except of $D_{|C||B|-1}^{(2)}$ (the values of ϕ for $u \in D_{|C||B|-1}^{(2)}$ are settled in the following steps and depend successively on b^{k+K}, b^{k+K+1}, etc., this will not be relevant for our purposes). So for the top level $D_{|B|-1}^{(1)} = \bigcup_{j=1}^{|C|} D_{j|B|-1}^{(2)}$ we set $\check{C} + b_{|B|-1}$ as the consecutive values of ϕ (remembering that $b_0 = 0$), that is

$$\phi|_{D_{j|B|-1}^{(2)}} = c_{j-1} + c_j + b_{|B|-1} \text{ for } j = 1, \ldots, |C| - 1. \tag{25}$$

We now define a (partial) function $f_{s,C}$ in the following way:

$$f_{s,C}|_{D_{j|B|-1}^{(2)}} = \check{c}_{j-1} + \check{c}_j + \ldots + \check{c}_{j+s-1} + sb_{|B|-1} + s\Sigma$$

for $j = 1, \ldots, |C| - s$ and then we spread this constant value down the column for which $D_{j|B|-1}^{(2)}$ is the top level. It follows that the function $f_{s,C}$ is not defined on the last s columns of the tower $\mathcal{D}^{(1)}$, that is, on a set of measure $s/|C|$. Moreover, the value

$$\big| \textstyle\int_Y (-1)^{f_{s,C}(u)} \, d\nu(u) \big| \text{ (defined up to } \pm s/|C|\text{) does not depend on } B \tag{26}$$

and

$$\Big| \big| \int_Y (-1)^{\phi^{(s|B||C|)}(u)} \, d\nu(u) \big| - \big| \int_Y (-1)^{f_{s,C}(u)} \, d\nu(u) \big| \Big| < s/|C|. \tag{27}$$

Come back now to the Thue-Morse sequence $x = 01 \times 01 \times \ldots$, use (9) and (10) and apply the above to obtain

$$\Big| \big| \int_Y (-1)^{\phi^{(s \cdot 2^{k+K})}(u)} \, d\nu(u) \big| - \big| \int_Y (-1)^{f_{s,C}(u)} \, d\nu(u) \big| \Big| < s/2^K.$$

Taking into account that $\widehat{\sigma}(s) = \widehat{\sigma}(s \cdot 2^\ell)$, as a conclusion of this, we obtain that for the Thue-Morse sequence we have

$$\Big| \int_Y (-1)^{f_{s,C}} \, d\nu \Big| \xrightarrow[|C| \to \infty]{} |\widehat{\sigma}(s)|. \tag{28}$$

Now, we borrow the idea of stabilizing subsequence from [1]. We say that a (generalized) Morse sequence $y = b^0 \times b^1 \times \ldots$ has a *stabilizing Thue-Morse subsequence* if there exists a subsequence $k_1 < k_2 < \ldots$ such that for each $K \geq 1$

$$(b^{k_i}, b^{k_i+1}, \ldots, b^{k_i+K-1}) = (01, 01, \ldots, 01) \ \text{ eventually in } i.$$

Denote by η the spectral measure of y. Using (10), (26), (27) and (28) we obtain that for each $i \geq 1$ and $K \geq 1$ (sufficiently large)

$$\big| |\widehat{\eta}(sq_{Kk_i})| - |\widehat{\sigma}(s)| \big| < s/2^K.$$

Assume that $r \neq s$ are odd numbers. Then, in view of Lemma 6, without loss of generality we can assume that $|\widehat{\sigma}(r)| \neq |\widehat{\sigma}(s)|$. By the above, we can indicate easily a subsequence (L_i) such that

$$|\widehat{\eta}(sq_{L_i})| \to |\widehat{\sigma}(s)| \ \text{ and } \ |\widehat{\eta}(rq_{L_i})| \to |\widehat{\sigma}(r)|.$$

Applying Corollary 1, we obtain $\widetilde{\eta}^{(s)} \perp \widetilde{\eta}^{(r)}$ and, moreover, Remark 2 applies for y. In view of Proposition 4, we have proved the following result.

Corollary 4. *Assume that $y = b^0 \times b^1 \times \ldots$ is a generalized Morse sequence possessing a stabilizing Thue-Morse subsequence. Then Sarnak's conjecture holds for the corresponding Morse dynamical system.*

References

[1] H. El Abdalaoui, M. Lemańczyk, T. de la Rue, *On spectral disjointness of powers for rank-one transformations and Möbius orthogonality*, arXiv:1301.0134.

[2] J. Bourgain, *On the correlation of the Möbius function with random rank-one systems*, Journal d'Analyse Math. (to appear), arXiv:1112.1032.

[3] J. Bourgain, P. Sarnak, T. Ziegler, *Disjointness of Möbius from horocycle flows*, in "From Fourier and Number Theory to Radon Transforms and Geometry", in memory of Leon Ehrenpreiss, Developments in Mathematics, **28** (2012), 67–83, Springer Verlag. arXiv:1110.0992.

[4] C. Dartyge, G. Tenenbaum, *Sommes des chiffres de multiples d'entiers*, Ann. Inst. Fourier (Grenoble) **55** (2005), 2423–2474.

[5] E. Glasner, *Ergodic Theory via Joinings*, Mathematical Surveys and Monographs **101**, AMS, Providence, RI, 2003.

[6] B. Green, T. Tao, *The Möbius function is strongly orthogonal to nilsequences*, Annals Math. **175** (2012), 541-566.

[7] M. Guenais, *Morse cocycles and simple Lebesgue spectrum*, Ergodic Theory Dynam. Systems **19** (1999), 437Ŭ446.

[8] M. Hochman, *Geometric rigidity of ×m invariant measures*, arXiv:1008.3548v2.

[9] K.-H. Indlekofer, I. Katai, *Investigations in the theory of q-additive and q-multiplicative functions.I*, Acta Math. Hungar. **91** (2001), 53-78.

[10] K. Jacobs, M. Keane, *0-1 sequnces of Toeplitz type*, Z. Wahr. Verw. Geb. **13** (1969), 123-131.

[11] A. del Junco, *A transformation with simple spectrum which is not rank one*, Canad. J. Math. **29** (1977), 655–663.

[12] A. Katok, J.-P. Thouvenot, *Spectral Properties and Combinatorial Constructions in Ergodic Theory*, Handbook of dynamical systems. Vol. 1B, 649–743, Elsevier B. V., Amsterdam, 2006.

[13] M. Keane, *Generalized Morse sequences*, Z. Wahr. Verw. Geb. **10** (1968), 335-353.

[14] J. Kwiatkowski, *Spectral isomorphism of Morse dynamical systems*, Bull. Pol. Acad. Sc. **29** (1981), 105-114.

[15] M. Lemańczyk, *Toeplitz $\mathbb{Z}_2$-extensions*, Annals H. Poincaré **24** (1988), 1-43.

[16] Y. Liu, P. Sarnak, *The Möbius function and distal flows*, arXiv:1303.4957

[17] M. Quefelec, *Substitution Dynamical Systems - Spectral Analysis*, Lecture Notes in Math. **1294** (1987).

[18] P. Sarnak, *Three lectures on the Möbius function, randomness and dynamics*, publications.ias.edu/sarnak/.

[19] T. Tao, http://terrytao.wordpress.com/2012/10/14/the-chowla-conjecture-and-the-sarnak-conjecture/

[20] T. Tao, http://mathoverflow.net/questions/97261/mobius-randomness-of-the-rudin-shapiro-sequence

[21] S. Williams, *Toeplitz minimal flows which are not uniquely ergodic*, Z. Wahr. Verw. Geb. **67** (1984), 95-107.

El Houcein El Abdalaoui:
Laboratoire de Mathématiques Raphaël Salem,
Université de Rouen, CNRS – Avenue de l'Université – 76801 Saint Étienne du Rouvray, France
elhoucein.elabdalaoui@univ-rouen.fr

Stanisław Kasjan, Mariusz Lemańczyk:
Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, 12/18 Chopin street, 87-100 Toruń, Poland
skasjan@mat.umk.pl
mlem@mat.umk.pl